Journal of Immunological Methods, 133 (1990) 297–298
Elsevier

Contents to volume 133 (1990)

Number 1 – 4 October 1990

Announcement from the publisher v

Research reports

A novel covalent enzyme-linked immunoassay (CELIA) for simultaneously measuring free and immune complex bound antibodies of defined specificity. I. Application to naturally occurring antipolyamine antibodies in human sera
A.M. Roch, J.G. Delcros, J.P. Ripoll, V. Thomas, J. Richard and G. Quash (France) 1

A novel covalent enzyme-linked immunoassay (CELIA) for simultaneously measuring free and immune complex bound antibodies with a defined specificity. II. Application to immune complexes containing viral antigens in human sera
V. Thomas, S. El Alaoui, A.M. Roch and G. Quash (France) 13

A turbidimetric assay in an ELISA reader for the determination of mononuclear phagocyte procoagulant activity
T.W. Jungi (Switzerland) 21

Isolation of rat peritoneal mononuclear and polymorphonuclear leucocytes on discontinuous gradients of Nycodenz
S. Fisker, K. Kudahl and O. Sonne (Denmark) 31

Fluorescein-specific hybridomas derived from primary mice exhibit more stringent growth requirements than do hybrids from pre-immune animals
J.B. Skeath, M.R. Dalesandro, M.A. Ferguson, G.C. Kinnel and J.A. Owen (U.S.A.) 39

A new method for measuring eosinophil activating factors, based on the increased expression of CR3 α chain (CD11b) on the surface of activated eosinophils
K.J.I. Thorne, B.A. Richardson, G. Mazza and A.E. Butterworth (U.K.) 47

Use of the fluorescence activated cell sorter to enrich dendritic cells from mouse spleen
M.T. Crowley, K. Inaba, M.D. Witmer-Pack, S. Gezelter and R.M. Steinman (U.S.A.) 55

Immobilization of C3b with retention of functional activity
R.E. Edens, R.J. Linhardt and J.M. Weiler (U.S.A.) 67

Induction of anti-idiotypic antibodies to a myeloma protein linked covalently to muramyl dipeptide
A.E. Eggers and J. Michl (U.S.A.) 71

Transfection and gene expression in normal and malignant primary B lymphocytes
M. Buschle, M.K. Brenner, I.S.Y. Chen, H.G. Drexler, S.M. Gignac and C.M. Rooney (U.K., U.S.A.) 77

New fluorescent dyes for lymphocyte migration studies. Analysis by flow cytometry and fluorescence microscopy
S.A. Weston and C.R. Parish (Australia) 87

Fetal bovine serum contains an inhibitor of interleukin-1
R.C. McKenzie, C.B. Harley, S. Matic and D.N. Sauder (Canada) 99

Characterization of polyclonal anti-peptide antibodies specific for transforming growth factor β_2
A.J.M. Van den Eijnden-Van Raaij, I. Koornneef, H.G. Slager, C.L. Mummery and E.J.J. Van Zoelen (The Netherlands) 107

Time resolved fluoroimmunoassay of 7-methyl-2′-deoxyguanosine imidazole (ring open)
P. Degan, R. Montesano, A. Abbondandolo and G. Montagnoli (Italy, France) 119

Bioassay of interleukin-1 in serum and plasma following removal of inhibitory activity with polyethylene glycol
S.J. Hopkins and M. Humphreys (U.K.) 127

A single-step method for the purification of anti-FITC antibodies by use of a coumarin immunosorbent
D. Samuel and R.A. Abuknesha (U.K.) 133

Short communication

Dissociation rate constant of the biotin-streptavidin complex
U. Piran and W.J. Riordan (U.S.A.) 141

Letter to the editors

Non-specific reactivity of monoclonal antibodies in ELISA due to the presence of an antigen-like structure on the alkaline phosphatase molecule
M. St. Laurent, F. Martel, S. Verrette and R. Lemieux (Canada) 145

Book reviews 147

Notes to authors 149

0022-1759/90/$03.50 © 1990 Elsevier Science Publishers B.V. (Biomedical Division)

Number 2 – 19 October 1990

Research reports

The sensitivity of dot immunoassay for the peptides helodermin, histidine-isoleucinamide (PHI) and histidine-methioninamide (PHM) increases after peptide cross-linking to proteins prefixed on nitrocellulose
P. Gourlet, M. Svoboda, A. Cauwin and J. Christophe (Belgium) 151

Antigen protection of monoclonal antibodies undergoing labelling
M. Ramjeesingh, M. Zywulko, A. Rothstein, R. Whyte and E.Y. Shami (Canada) 159

An immunoassay for the quantitation of cell membrane MHC class II antigen
C.V. Whiting and P.W. Bland (U.K.) 169

The use of diphenylene iodonium, an inhibitor of NADPH oxidase, to investigate the antimicrobial action of human monocyte derived macrophages
A.K. Robertson, A.R. Cross, O.T.G. Jones and P.W. Andrew (U.K.) 175

Quantitation of components of the alternative pathway of complement (APC) by enzyme-linked immunosorbent assays
M. Oppermann, H. Baumgarten, E. Brandt, W. Gottsleben, C. Kurts and O. Götze (F.R.G.) 181

A chemiluminescent, microparticle-membrane capture immunoassay for the detection of antibody to hepatitis B core antigen
J. Wolf-Rogers, J.A. Weare, K. Rice, E.F. Robertson, P. Guidinger, O.S. Khalil and G. Madsen (U.S.A.) 191

Functional assay of C5-activating and nonactivating cobra venom factor preparations in the mouse system
C.W. Van den Berg, P.C. Aerts and H. Van Dijk (The Netherlands) 199

Effect of hapten heterology on thyroid hormone immunoassays
U. Piran, W.J. Riordan and D.R. Silbert (U.S.A.) 207

The detection of anti-Ro/SS-A and anti-La/SS-B antibodies. A comparison of counterimmunoelectrophoresis with immunoblot, ELISA, and RNA-precipitation assays
J.F. Meilof, I. Bantjes, J. De Jong, A.P. Van Dam and R.J.T. Smeenk (The Netherlands) 215

A comparison of immunoblotting, flow cytometry and ELISA to monitor the binding of anti-lipopolysaccharide monoclonal antibodies
D. Nelson, W. Neill and I.R. Poxton (U.K.) 227

Gating of the so-called 'lymphocytic' cell population for the quantification of natural killer cells ($CD16^+$) by flow cytometry causes loss of CD16 positive cells
A.M. Lagaay, C.B. Bosman, M. Van der Keur, G.J. Ligthart, H.R.E. Schuit, H.J. Tanke and W. Hijmans (The Netherlands) 235

A method for blocking antigen-independent binding of human IgM to frozen tissue sections when screening human hybridoma antibodies
H. Ditzel, K. Erb, B. Nielsen, P. Borup-Christensen and J.C. Jensenius (Denmark) 245

Direct blotting with viable cells of protein mixtures separated by two-dimensional gel electrophoresis
H. Gulle, B. Schoel and S.H.E. Kaufmann (F.R.G.) 253

Qualitative analysis of antibody binding. An in vitro assay for the evaluation and development of vaccines
U. Bruderer, E. Fürer, S.J. Cryz, Jr. and A.B. Lang (Switzerland) 263

Purification of human basophils using mouse monoclonal IgE
J.T. Schroeder and L.R. Hanrahan, Jr. (U.S.A.) 269

An enzyme-linked immunosorbent assay for the measurement of human interleukin-6
N. Ida, S. Sakurai, T. Hosaka, K. Hosoi, T. Kunitomo, Y. Matsuura and M. Kohase (Japan) 279

A rapid [^{3}H]glycerol radioassay for determination of monocyte-mediated growth inhibition of *Mycobacterium avium*
D.K. Blanchard, M.B. Michelini-Norris and J.Y. Djeu (U.S.A.) 285

Editorial 291

Book review 293

Announcements 295

Contents to volume 133 (1990) 297

Notes to authors 299